

DISCRETE COMPONENTS OF SOME COMPLEMENTARY SERIES

B.SPEH AND T. N. VENKATARAMANA

ABSTRACT. We show that Complementary Series representations of $SO(n, 1)$, which are sufficiently close to a cohomological representation contain **discretely**, Complementary Series of $SO(m, 1)$ also sufficiently close to cohomological representations, provided that the degree of the cohomological representation does not exceed $m/2$.

We prove, as a consequence, that the cohomological representation of degree i of the group $SO(n, 1)$ contains discretely, the cohomological representation of degree i of the subgroup $SO(m, 1)$ if $i \leq m/2$.

As a global application, we show that if $G/\mathbb{Q}$ is a semisimple algebraic group such that $G(\mathbb{R}) = SO(n, 1)$ up to compact factors, and if we assume that for all n , the tempered cohomological representations are not limits of complementary series *in the automorphic dual of $SO(n, 1)$* , then for all n , non-tempered cohomological representations are isolated in the automorphic dual of G . This reduces conjectures of Bergeron to the case of **tempered** cohomological representations.

1991 *Mathematics Subject Classification*. Primary 11F75; Secondary 22E40, 22E41

B. Speh, Department of Mathematics, 310, Malott Hall, Cornell University, Ithaca, NY 14853-4201, U.S.A.

T. N. Venkataramana, School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay - 400 005, INDIA. venky@math.tifr.res.in.

1. INTRODUCTION

A famous Theorem of Selberg [Sel] says, that the non-zero eigenvalues of the Laplacian acting on functions on quotients of the upper half plane by *congruence subgroups* of the integral modular group are bounded away from zero, as the congruence subgroup varies. In fact, Selberg proved that every non-zero eigenvalue λ of the Laplacian on functions on $\Gamma \backslash \mathfrak{h}$, $\Gamma \subset SL_2(\mathbb{Z})$, satisfies the inequality

$$\lambda \geq \frac{3}{16}.$$

A Theorem of Clozel [Clo] generalises this result to any congruence quotient of any symmetric space of non-compact type: if G is a linear semi-simple group defined over $\mathbb{Q}$, $\Gamma \subset G(\mathbb{Z})$ a congruence subgroup and $X = G(\mathbb{R})/K$ the symmetric space of G , then nonzero eigenvalues λ of the Laplacian *acting on the space of functions on $\Gamma \backslash X$* satisfy:

$$\lambda \geq \epsilon,$$

where $\epsilon > 0$ is a number **independent** of the congruence subgroup Γ .

Analogous questions on Laplacians acting on differential forms of higher degree (functions may be thought of as differential forms of degree zero) have geometric implications on the cohomology of the locally symmetric space. Concerning the eigenvalues of the Laplacian, Bergeron ([Ber]) has conjectured the following:

Conjecture 1. (Bergeron) Let X be the real hyperbolic n -space and $\Gamma \subset SO(n, 1)$ a congruence arithmetic subgroup. The non-zero eigenvalues λ of the Laplacian acting on the space $\Omega^i(\Gamma \backslash X)$ of differential forms of degree i satisfy:

$$\lambda \geq \epsilon,$$

for some $\epsilon > 0$ **independent** of the congruence subgroup Γ , provided i is strictly less than the “middle dimension” (i.e. $i < [n/2]$).

If n is even, he conjectures that the above conclusion holds even if $i = [n/2] = n/2$. (When n is odd, there is a slightly more technical statement which we omit).

In this paper, we show, for example, that if the above conjecture holds true in the *middle degree* for all even integers n , then the conjecture holds for arbitrary degrees of the differential forms (See Theorem 3). For odd n , there is, again, a more technical statement (see Theorem 3, part (2)).

The main Theorem of the present paper is Theorem 1 on the occurrence of some discrete components of certain Complementary Series Representations associated to $SO(n, 1)$. The statement on Laplacians may be deduced from the main theorem, using the Burger-Sarnak ([Bu-Sa]) method.

We now describe the main theorem more precisely. Let $i \leq [n/2] - 1$ and $G = SO(n, 1)$. Let $P = MAN$ be the Langlands decomposition of P , $K \subset G$ a maximal compact subgroup of G containing M . Then K is isomorphic to the orthogonal group $O(n)$ and $M = O(n - 1)$; denote by $\mathfrak{p}$ the standard representation of $O(n)$, tensored by the determinant character; if $\mathfrak{g}, \mathfrak{k}$ denote the complexified Lie algebras of G and K , then the Cartan decomposition takes the form $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, as K -modules.

Let, analogously, $\mathfrak{p}_M$ be the standard representation of M twisted by the character such that this representation occurs in the restriction of $\mathfrak{p}$ to the subgroup M of K . Let $\wedge^i$ be its i -th exterior power representation. Denote by ρ_P^2 the character of P acting on the top exterior power of the Lie algebra of N . Consider the induced representation (with normalized induction)

$$\widehat{\pi_u(i)} = \text{Ind}_P^G(\wedge^i \otimes \rho_P(a)^u)$$

for $0 < u < 1 - \frac{2i}{n-1}$. The representation $\widehat{\pi_u(i)}$ denotes the **completion** of the space $\pi_u(i)$ of K -finite functions with respect to the G -invariant metric on $\pi_u(i)$, and is called the **complementary series representation** corresponding to the representation $\wedge^i$ of M and the parameter u .

Let $H = SO(n - 1, 1)$ be embedded in G such that $P \cap H$ is a maximal parabolic subgroup of H , $A \subset H$, and $K \cap H$ a maximal compact subgroup of H . We now assume that $\frac{1}{n-1} < u < 1 - \frac{2i}{n-1}$. Let $u' = \frac{(n-1)u-1}{n-2}$; then $0 < u' < 1 - \frac{2i}{n-2}$. Denote by $\wedge_H^i$ the i -th exterior power of the standard representation of $M \cap H \simeq O(n - 2)$.

We obtain analogously the complementary series representation

$$\widehat{\sigma_{u'}(i)} = \text{Ind}_{P \cap H}^H(\wedge_H^i \otimes \rho_{P \cap H}(a)^{u'}),$$

of H . The main theorem of the paper is the following.

Theorem 1. *If*

$$\frac{1}{n-1} < u < 1 - \frac{2i}{n-1},$$

then the complementary series representation $\widehat{\sigma_u(i)}$ occurs discretely in the restriction of the complementary series representation $\widehat{\pi_u(i)}$ of $G = SO(n, 1)$, to the subgroup $H = SO(n - 1, 1)$:

$$\widehat{\sigma_u(i)} \subset \widehat{\pi_u(i)}|_{SO(n-1,1)}.$$

Remark 1. The corresponding statement is false for the space of K -finite vectors in both the spaces; this inclusion holds only at the level of completions of the representations involved.

As u tends to the limit $1 - \frac{2i}{n-1}$, the representations $\widehat{\pi_u(i)}$ tend (in the Fell topology on the unitary dual $\widehat{G}$) both to the representation $\widehat{A_i} = \widehat{A_i(n)}$ and to $\widehat{A_{i+1}} = \widehat{A_{i+1}(n)}$; we denote by $\widehat{A_j(n)}$ the unique cohomological representation of G which has cohomology (with trivial coefficients) in degree j and by $A_j(n)$ the space of K -finite vectors in $\widehat{A_j(n)}$. Using this, and the proof of Theorem 1, we obtain

Theorem 2. *The restriction of the cohomological representation $\widehat{A_i(n)}$ of $SO(n, 1)$ to the subgroup $H = SO(n - 1, 1)$ contains discretely, the cohomological representation $\widehat{A_i(n - 1)}$ of $SO(n - 1, 1)$:*

$$\widehat{A_i(n - 1)} \subset \widehat{A_i(n)}|_{SO(n-1,1)}.$$

Suppose now that G is a semi-simple linear algebraic group defined over $\mathbb{Q}$ and $\mathbb{Q}$ -simple, such that

$$G(\mathbb{R}) \simeq SO(n, 1)$$

up to compact factors with $n \geq 4$ (if $n = 7$, we assume in addition that G is not the inner form of some triality D_4). Then there exists a $\mathbb{Q}$ -simple $\mathbb{Q}$ -subgroup $H_1 \subset G$ such that

$$H_1(\mathbb{R}) \simeq SO(n - 2, 1),$$

up to compact factors.

Denote by $\widehat{G}_{Aut}$ the “automorphic dual” of $SO(n, 1)$, in the sense of Burger-Sarnak ([Bu-Sa]). Suppose $\widehat{A_i} = \widehat{A_i(n)}$ is a limit of representations $\widehat{\rho_m}$ in $\widehat{G}_{Aut}$. The structure of the unitary dual of $SO(n, 1)$ shows that this means that $\widehat{\rho_m} = \widehat{\pi_{u_m}(i)}$ for some sequence $\widehat{u_m}$ which tends from the left, to $1 - \frac{2i}{n-1}$ (or to $1 - \frac{2i+2}{n-1}$; we will concentrate only on the first case, for ease of exposition).

Since $\widehat{\rho_m} = \widehat{\pi_{u_m}(i)} \in \widehat{G}_{Aut}$, a result of Burger-Sarnak ([Bu-Sa]), implies that

$$res(\widehat{\rho_m}) \subset \widehat{H}_{Aut}.$$

Applying Theorem 1 twice, we get

$$\widehat{\sigma}_m = \widehat{\sigma}_{u''_m}(i) \in \text{res}(\widehat{\rho}_m) \subset \widehat{H}_{Aut}$$

Taking limits as m tends to infinity, we get $\widehat{A_i(n-2)}$ as a limit of representations $\widehat{\sigma}_m$ in $\widehat{H}_{Aut}$. Therefore, the isolation of $\widehat{A_i(n)}$ in $\widehat{G}_{Aut}$ is reduced to that for $SO(n-2, 1)$, and so on. We can finally assume that $\widehat{A_i(m)}$ is a tempered representation of $SO(m, 1)$ where $m = 2i$ or $2i + 1$.

This proves the following Theorem.

Theorem 3. *If for all m , the tempered cohomological representation $\widehat{A_i(m)}$ (i.e. when $i = [m/2]$) is not a limit of complementary series in the automorphic dual of $SO(m, 1)$, then for all integers n , and for $i < [n/2]$, the cohomological representation $\widehat{A_i(n)}$ is isolated in the automorphic dual of $SO(n, 1)$.*

The major part of the paper is devoted to proving the main theorem (Theorem 1). The proof of Theorem 1 proceeds as follows.

(1) We first prove Theorem 1 when $i = 0$; that is $\widehat{\pi}_u = \widehat{\pi}_u(0)$ is an unramified representation and $\frac{1}{n-1} < u < 1$. In this case, we get a model of the representation $\widehat{\pi}_u$ by restricting the functions (sections of a line bundle) on G/P to the big Bruhat cell $Nw \simeq \mathbb{R}^{n-1}$ and taking their Fourier transforms. The G -invariant metric is particularly easy to work with on the Fourier transforms (see Theorem 10); it is then easy to see that there is an isometric embedding $J : \widehat{\sigma}_{u'} \rightarrow \widehat{\pi}_u$ (Theorem 15).

(2) We now describe the proof of Theorem 1 in the ramified case. The representation $\widehat{\pi}_u(i)$ is the “completion” of the space of Schwartz class functions on the big Bruhat cell, with values in the representation $\wedge^i$ of the compact group M , with respect to a bilinear form. That this bilinear form is an inner product is not so clear (in contrast to the unramified case); we assume this ([Ba-B]; in [Ba-B], they prove the existence of a positive definite inner product on the space of K -finite vectors in $\widehat{\pi}_u(i)$; since our bilinear form is also G -invariant, the irreducibility of the module involved shows that these two bilinear forms coincide.

We again get a model of the representations $\widehat{\pi}_u(i)$ on the space of Fourier transforms of Schwartz class functions on the big Bruhat Cell in G/P . There is an embedding $J : \widehat{\sigma}_{u'}(i) \rightarrow \widehat{\pi}_u(i)$, which can be explicitly described. We prove that the adjoint J^* of J is essentially the

restriction $\pi_{-u}(i) \rightarrow \sigma_{-u}(i)$. This proves that the map J is equivariant for the action of the subgroup H and establishes the main theorem (Theorem 1) for arbitrary i .

2. PRELIMINARY RESULTS AND NOTATION

2.1. The Group $G = SO(n, 1)$ and the representations $\pi_u(i)$.
 The group $G = SO(n, 1)$ is the subgroup of $SL_{n+1}(\mathbb{R})$ preserving the quadratic form

$$x_0x_n + x_1^2 + x_2^2 + \cdots + x_{n-1}^2,$$

on $\mathbb{R}^{n+1}$. Denote by $e_0, e_1, \dots, e_{n-1}, e_n$ the standard basis of $\mathbb{R}^{n+1}$. Denote by w the matrix in $SL_{n+1}(\mathbb{R})$, which takes the basis element e_0 to e_n , e_n to $-e_0$ and takes the other basis elements e_i into e_i for $1 \leq i \leq n-1$. Then w has the matrix form

$$\begin{pmatrix} 0 & 0_{n-1} & 1 \\ {}^t0 & 1_{n-1} & 0 \\ -1 & 0_{n-1} & 0 \end{pmatrix}$$

where 0_{n-1} is the zero vector in $\mathbb{R}^{n-1}$ viewed as a row vector of size $n-1$ and t0 refers to its transpose (a column vector of size $n-1$ all of whose entries are zero) and 1_{n-1} refers to the identity matrix of size $n-1$. The group G is the matrix group

$$\{g \in GL_{n+1}(\mathbb{R}) : {}^t wg = w\}.$$

Denote by P the subgroup of G which takes the basis vector e_0 into a multiple of itself. Let A be the group of diagonal matrices in G , M the maximal compact subgroup of the centralizer of A in G . Let N be the unipotent radical of P . Then,

$$A = \{d(a) = \begin{pmatrix} a & 0_{n-1} & 0 \\ {}^t0 & 1_{n-1} & {}^t0 \\ 0 & 0_{n-1} & a^{-1} \end{pmatrix} : a \in \mathbb{R}^*\},$$

Similarly M consists of matrices of the form

$$M = \left\{ \begin{pmatrix} \pm 1 & 0_{n-1} & 0 \\ {}^t0 & m & {}^t0 \\ 0 & 0_{n-1} & \pm 1 \end{pmatrix} : m \in SO(n-1) \right\}.$$

The group N is

$$N = \{u(x) = \begin{pmatrix} 1 & x & \frac{|x|^2}{2} \\ {}^t0 & 1_{n-1} & {}^tx \\ 0 & 0_{n-1} & 1 \end{pmatrix} : x \in \mathbb{R}^{n-1}\},$$

where $x \in \mathbb{R}^{n-1}$ is thought of as a row vector. Then $P = MAN$ is the Langlands decomposition of P . The map $x \mapsto u(x)$ from $\mathbb{R}^{n-1}$ into N is an isomorphism of algebraic groups.

Let $K = G \cap O(n+1)$. Then K is a maximal compact subgroup of G and is isomorphic to $O(n) \times O(1)$. Let $\rho_P(a)$ be the character whose square is the determinant of $Ad(d(a))$ acting on the complex Lie algebra $\mathfrak{n} = Lie(N) \otimes \mathbb{C}$. Then $\rho_P(a) = a^{\frac{n-1}{2}}$.

Write $\mathfrak{g} = Lie(G) \otimes \mathbb{C}$, $\mathfrak{k} = Lie(K) \otimes \mathbb{C}$ and set $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, the Cartan Decomposition which is complexified. As a representation of $K = O(n)$, the space $\mathfrak{p} = \mathbb{C}^n$ is the standard representation twisted by a character. Denote by $\mathfrak{p}_M = \mathbb{C}^{n-1}$ the standard representation of $M = O(n-1)$ twisted by that character so that $\mathfrak{p}_M$ occurs in $\mathfrak{p}$ restricted to K . Let i be an integer with $0 \leq i \leq [n/2] - 1$ where $[]$ denotes the integral part. Denote by $\wedge^i \mathfrak{p}_M$ the representation of M on the i -th exterior power of $\mathfrak{p}_M$.

If $u \in \mathbb{C}$ and $p = man \in MAN = P$, the map

$$p \mapsto \wedge^i \mathfrak{p}_M(m) \otimes \rho_P(a)^u \otimes \text{triv}_N(n) = \wedge^i(m) \rho_P(a)^u$$

is a representation of P . Define

$$\pi_u(i) = \text{Ind}_P^G(\wedge^i \mathfrak{p}_M \otimes \rho_P(a)^u \otimes 1_N)$$

to be the space of functions on G with values in the vector space $\wedge^i \mathfrak{p}_M$ which satisfy

$$f(gman) = \wedge^i(m)(f(g))\rho_P(a)^{-(1+u)},$$

for all $(g, m, a, n) \in G \times M \times A \times N$. In addition, the function f is assumed to be K -finite. The space $\pi_u(i)$ is a $(\mathfrak{g}, K)$ -module. The representation $\pi_u(i)$ is called the complementary series of $SO(n, 1)$ with parameter u and $\wedge^i$.

When $i = 0$, the representation $\pi_u(0)$ has a K -fixed vector, namely the function which satisfies $f(kan) = \rho_P(a)^{-(1+u)}$ for all $kan \in KAN = G$. We denote $\pi_u(0) = \pi_u$; we refer to this representation as the unramified complementary series of $SO(n, 1)$ with parameter u .

2.2. Bruhat Decomposition of $SO(n, 1)$. We have the **Bruhat Decomposition** of $O(n, 1)$:

$$(1) \quad G = NwMAN \cup MAN.$$

That is, every matrix $g \in G = O(n, 1)$ which does not lie in the parabolic subgroup P , may be written in the form $g = u(x)wmd(a')u(y)$ for some $x, y \in \mathbb{R}^{n-1}$, $a' \in \mathbb{R}^*$ and $m \in M$ or in the form $g = md(a)u(y)$. In these coordinates, the set $NwP = NwMAN$ is an open subset of

G of full Haar measure. The map $x \mapsto u(x)$ is an isomorphism of groups from $\mathbb{R}^{n-1}$ onto N . Let dn denote the image of the Lebesgue measure on $\mathbb{R}^{n-1}$ under this map. Denote by d^*a the measure $\frac{da'}{|a'|}$ for $a' \in \mathbb{R}^* \simeq A$ via the map $a' \mapsto d(a')$. Under the Bruhat decomposition (equation (1)), there exists a choice of Haar measure on M such that

$$(2) \quad dg = dx \, d^*a' \, dm \, dy \, \rho_P(a')^2.$$

Given $x = (x_1, x_2, \dots, x_k) \in \mathbb{R}^k$, denote by $|x|^2$ the sum

$$|x|^2 = \sum_{j=1}^k x_j^2.$$

Lemma 4. *If $x \in \mathbb{R}^{n-1}$ and $x \neq 0$, then the Bruhat decomposition of the element $g = wu(x)w$ is given by*

$$g = u(z)wm(x)d(a')u(y)$$

with

$$z = \frac{2x}{|x|^2}, \quad m = 1_{n-1} - 2 \frac{{}^t x x}{|x|^2}, \quad a' = \frac{|x|^2}{2} \text{ and } y = x$$

(note that ${}^t x x$ is a square matrix of size $n-1$ and that m is the reflection on the orthogonal complement of x).

The proof is by multiplying out the right hand side and explicitly computing the matrix product.

We will find it convenient to compute the Jacobian of the transformation $x \mapsto \frac{2x}{|x|^2}$ for $x \in \mathbb{R}^{n-1} \setminus \{0\}$.

Lemma 5. *If $f \in \mathcal{C}_c(\mathbb{R}^{n-1} \setminus \{0\})$ is a continuous function with compact support, then we have the formula*

$$\int_{\mathbb{R}^{n-1}} dx f\left(\frac{x}{|x|^2}\right) = \int_{\mathbb{R}^{n-1}} \frac{dx}{\left(\frac{|x|^2}{2}\right)^{n-1}} f(x).$$

Proof. In polar coordinates, $x = rp$ with $r > 0$ and p in the unit sphere S^{n-2} of $\mathbb{R}^{n-1}$. Then the Lebesgue measure dx has the form $dx = r^{n-1} \frac{dr}{r} d\mu(p)$ where μ is a rotation invariant measure on S^{n-2} . In polar coordinates, the transformation $x \mapsto \frac{2x}{|x|^2}$ takes the form $(r, p) \mapsto (\frac{2}{r}, p)$. Therefore $d(\frac{x}{|x|^2}) = \frac{2^{n-1}}{r^{n-1}} \frac{dr}{r} d\mu(p) = \frac{dx}{r^{2n-2}/2^{n-1}}$. $\square$

2.3. Iwasawa Decomposition of $O(n, 1)$. We have the **Iwasawa decomposition** of $O(n, 1) = G$:

$$(3) \quad G = KAN.$$

That is, every matrix g in $SO(n, 1)$ may be written as a product $g = kd(a)u(y)$ with $k \in K$, $a \in \mathbb{R}^*$ and $y \in \mathbb{R}^{n-1}$.

Lemma 6. *The Iwasawa decomposition of the matrix $u(x)w$ is of the form*

$$u(x)w = k(x)d(a)u(y),$$

with $a = 1 + \frac{|x|^2}{2}$ and $y = \frac{x}{a} = \frac{x}{1 + \frac{|x|^2}{2}}$. The matrix $k(x)$ is given by the formula

$$k(x) = u(x)wu(-y)d(a^{-1}).$$

Proof. Put $g = u(x)w = kd(a)u(y)$. Compute the product

$$w^t u(x)u(x)w =^t gg =^t u(y)d(a)d(a)u(y).$$

By comparing the matrix entries on both sides of this equation, we get the Lemma. $\square$

We normalize the Haar measures dg , dk , d^*a and dy on G , K , A , N respectively so that

$$(4) \quad dg = \rho_P(a)^2 dk d^*a dy.$$

(dy is the Lebesgue measure on $N \simeq \mathbb{R}^{n-1}$, on $A \simeq \mathbb{R}^*$ the measure is $d^*a = \frac{d_1(a)}{|a|}$ where d_1a is the Lebesgue measure).

2.4. The space $\mathcal{F}_1$ and a linear form on $\mathcal{F}_1$. Define $\mathcal{F}_1$ to be the space of complex valued continuous functions f on G such for all $(g, m, a, n) \in G \times M \times A \times N$ we have

$$f(gman) = \rho_P(a)^{-2} f(g).$$

The group G acts on $\mathcal{F}_1$ by left translations. Denote by $\mathcal{C}_c(G)$ the space of complex valued continuous functions with compact support. The following result is well known.

Lemma 7. *The map*

$$f \mapsto \overline{f}(g) \stackrel{\text{def}}{=} \int_{MAN} f(gman) \rho_P(a)^2 dm dn d^*a,$$

is a surjection from $\mathcal{C}_c(G)$ onto the space $\mathcal{F}_1$.

It follows from this and the Iwasawa decomposition (see equation (4)), that the map

$$(5) \quad L(\bar{f}) = \int_K f(k) dk$$

is a G -invariant linear form on the space $\mathcal{F}_1$.

Lemma 8. *For every function $\phi \in \mathcal{F}_1$ we have the formula*

$$L(\phi) = \int_{\mathbb{R}^{n-1}} dx \phi(u(x)w) = \int_K dk \phi(k).$$

Proof. Given $f \in \mathcal{C}_c(G)$, we have from equation (2) that

$$\int_{\mathbb{R}^{n-1}} dx \bar{f}(u(x)w) = \int_G f(g) dg = L(\bar{f}).$$

On the other hand, by equation (4), we have

$$\int_K dk \bar{f}(k) = \int_G f(g) dg = L(\bar{f}).$$

These two equations show that the lemma holds if $\phi = \bar{f}$.

It follows from Lemma 7 that every function $\phi \in \mathcal{F}_1$ is of the form $\bar{f}$ for some f . This proves the Lemma. □

3. UNRAMIFIED COMPLEMENTARY SERIES FOR $SO(n, 1)$

3.1. A Model for Unramified Complementary Series. Let $0 < u < 1$, and $W = \mathbb{R}^{n-1}$. Denote by $\mathcal{S}(W)$ the Schwartz Space of W .

Let $K = O(n)$ and $M = O(n-1)$. Then K/M is the unit sphere in W . Denote by w the non-trivial Weyl group element of $SO(n, 1)$. The space G/P is homogeneous for the action of K and the isotropy at P is M . Therefore, $G/P = K/M$. The Bruhat decomposition implies that we have an embedding $N \subset K/M$ via the map $n \mapsto nwP \subset G/P = K/M$. Denote by $\mathcal{F}_u$ the space of **continuous** functions on G which transform according to the formula

$$f(gman) = f(g)\rho_P(a)^{-(1+u)},$$

for all $g \in G, m \in M, a \in A$ and $n \in N$. An element of the Schwartz space $\mathcal{S}(W)$ on $W \simeq N$ may be extended to a continuous function on G , which lies in the space $\mathcal{F}_u$.

We will view $\mathcal{S}(W)$ as a subspace of $\mathcal{F}_u$ via this identification.

Lemma 9. *If $0 < u < 1$ and $\phi \in \mathcal{S}(W) \subset \mathcal{F}_u$, the integral*

$$I_u(\phi)(g) = \int_W dx \phi(gu(x)w)$$

converges.

The integral $I(u)(\phi)$ lies in the space $\mathcal{F}_{-u}$.

Proof. The Iwasawa decomposition of $u(x)w$ (Lemma 6) implies that $u(x)w = kman$ with

$$\rho_P(a)^{1+u} = \frac{1}{(1 + \frac{|x|^2}{2})^{(n-1)(1+u)/2}}$$

where $|x|^2$ is the standard inner product of $n = u(x) \in N \simeq W$ in the space W and $k \in K, m \in M$ and $n \in N$. Therefore, for large $x \in W$, $\rho_P(a)^{1+u} \simeq |x|^{-(n-1)(1+u)}$, which proves the convergence of the integral. The last statement is routine to check. $\square$

There is a G -invariant pairing between $\mathcal{F}_u$ and $\mathcal{F}_{-u}$ defined by

$$\langle \phi, \psi \rangle = \int_K dk \overline{\phi(k)} \psi(k),$$

for all $\phi \in \mathcal{F}_u$ and $\psi \in \mathcal{F}_{-u}$. Here $\overline{\phi(k)}$ denotes the complex conjugate of $\phi(k)$.

Observe that the product $\overline{\phi(g)}\psi(g)$ of the elements $\phi \in \mathcal{F}_u$ and $\psi \in \mathcal{F}_{-u}$ lies in the space $\mathcal{F}_1$. Therefore, $\langle \phi, \psi \rangle = L(\phi\psi)$ where L is the invariant linear form on $\mathcal{F}_1$ as in equation (5).

From Lemma 8 it follows, for $\phi \in \mathcal{F}_u$ and $\psi \in \mathcal{F}_{-u}$, that

$$\langle \phi, \psi \rangle = \int_{\mathbb{R}^{n-1}} dx \overline{\psi(u(x)w)} \psi(u(x)w).$$

Given $\phi \in \mathcal{S}(W)$, denote by $\widehat{\phi}$ its Fourier transform.

Theorem 10. *For $0 < u < 1$ and $\phi \in \mathcal{S}(W) \subset \mathcal{F}_u$, we have the formula*

$$\langle \phi, I(u)(\phi) \rangle_u = c \int_W |\widehat{\phi}(x)|^2 \frac{1}{|x|^{(n-1)u}} dx,$$

relating the pairing between $\mathcal{F}_u$ and $\mathcal{F}_{-u}$, and a norm on the Schwartz space. (here c is a constant depending only on the constant u).

In particular, the pairing $\langle \phi, I(u)(\phi) \rangle$ is positive definite on the Schwartz space $\mathcal{S}(W)$.

The proof of Theorem 10 will occupy the rest of the section (3.1). Note that Theorem 10 is already proved in the paper of Grave-Viskik ([GV]). The proof we give below is a little different, and moreover, we need the functional equation that is proved below, in the course of the proof of Theorem 10.

Lemma 11. *If $\phi^* \in \mathcal{S}(W) \subset \mathcal{F}_u$, then we have the formula*

$$\int_N \phi^*(n'wnw)dn = \int_W \phi^*(y+x) \frac{1}{|x|^{(n-1)(1-u)}} dx,$$

where $n, n' \in N$ correspond respectively to $y, x \in W$.

Proof. We view $\mathcal{S}(W)$ as a subspace of $\mathcal{F}_u$. Thus, the function $\phi^*(x) \in \mathcal{S}(W)$ is identified with the section $x \mapsto \phi^*(u(x)w)$ with $\phi^* \in \mathcal{F}_u$. We then get the equality of the integrals:

$$\int_N \phi^*(nwn'w)dn' = \int_W \phi^*(u(y)wu(x)w)dx.$$

By Lemma 4,

$$wu(x)w = u\left(\frac{2x}{|x|^2}\right)m(x)d\left(\frac{|x|^2}{2}\right)u(x).$$

Using the fact that ϕ^* lies in $\mathcal{F}_u$, we see that

$$\phi^*(u(y)wu(x)w) = \phi^*\left(u\left(y + \frac{2x}{|x|^2}\right)\right)\left(\frac{|x|^2}{2}\right)^{-(n-1)(1+u)/2}.$$

Therefore, the integral of the lemma is

$$\int_W dx \phi^*(y + \frac{2x}{|x|^2}) (\frac{|x|^2}{2})^{-(n-1)(1+u)/2}.$$

We now use the computation of the Jacobian of the map $x \mapsto \frac{2x}{|x|^2}$ (see Lemma 5) to get

$$\int_N \phi^*(nwn'w) dn' = \int_W \frac{dx}{(|x|^2/2)^{n-1}} \phi^*(y+x) (\frac{|x|^2}{2})^{(n-1)(1+u)/2},$$

and this is easily seen to be the right hand side of the equation of the Lemma. $\square$

Recall that

$$I(u)(\phi^*)(nw) = \int_N dn' \phi^*(nwn'w).$$

From Lemma 8 we have

$$\langle \phi^*, I(u)(\phi^*) \rangle = \int_N dn (\phi^*(nw) I(u)(\phi^*)(nw)).$$

Therefore, it follows from the preceding Lemma that the G -invariant pairing $\langle \phi^*, I(u)(\phi^*) \rangle$ is the integral

$$\int_W dy \int_W dx \frac{1}{|x|^{(n-1)u}} \overline{\phi(y)} \phi(y+x).$$

The Fubini Theorem implies that this integral is

$$\int_W dx \frac{1}{|x|^{(n-1)u}} \int_W dy \overline{\phi(y)} \phi(y+x).$$

The inner integral is the L^2 inner product between ϕ and its translate by x . Since the Fourier transform preserves the inner product and converts translation by x into multiplication by the character $y \mapsto e^{-2ix \cdot y}$, it follows that the integral becomes

$$\int_W dx \frac{1}{|x|^{(n-1)(1-u)}} \int_W dy |\widehat{\phi}(y)|^2 e^{-2ix \cdot y}.$$

The integral over y is simply the Fourier transform. Therefore, we have

$$(6) \quad \langle \phi, I(u)(\phi) \rangle = \int_W dx \frac{1}{|x|^{(n-1)(1-u)}} |\widehat{\phi(y)}|^2(x).$$

Lemma 12. *Let $f \in \mathcal{S}(W)$, and s a complex number with real part positive and less than $(n-1)/2$. Then we have the functional equation*

$$\Gamma(s) \int_W dx \frac{1}{|x|^{2s}} \widehat{f}(x) = \Gamma(\frac{n-1}{2} - s) \int_W dx |x|^{n-1-2s} f(x),$$

where Γ is the classical Gamma function.

Proof. If s is a complex number with positive real part, then $\Gamma(s)$ is defined by the integral

$$\int_0^{+\infty} \frac{dt}{t} t^s e^{-t}.$$

Denote by $L(s, \widehat{f})$ the integral

$$\int_W dx \frac{1}{|x|^s} \widehat{f}(x).$$

Multiplying $L(2s, \widehat{f})$ by $\Gamma(s)$, and using Fubini, we obtain

$$\Gamma(s)L(2s, \widehat{f}) = \int_W \frac{dx}{|x|^{2s}} \widehat{f}(x) \int_0^\infty e^{-t} t^s \frac{dt}{t}.$$

Changing the variable t to $|x|^2 t$ and using Fubini again, we get

$$\Gamma(s)L(2s, \widehat{f}) = \int_0^\infty \frac{dt}{t} t^s \int_W e^{-t|x|^2} \widehat{f}(x) dx.$$

The inner integral is the inner product in $L^2(W)$ of the functions $e^{-t|x|^2}$ and $\widehat{f}$. The Fourier transform preserves this inner product. Therefore, the inner product is the inner product of their Fourier transforms.

The Fourier transform of $e^{-t|x|^2}$ is $\frac{1}{t^{(n-1)/2}} e^{-|x|^2/t}$. The Fourier transform of $\widehat{f}$ is $f(-x)$. Therefore, we have the equality

$$\Gamma(s)L(2s, \widehat{f}) = \int_0^\infty \frac{dt}{t} t^{s-(n-1)/2} \int_W e^{-|x|^2/t} f(-x) dx.$$

In this double integral, change t to $|x|^2/t$. We then get

$$\Gamma(s)L(2s, \widehat{f}) = \int_0^\infty \frac{dt}{t} t^{(n-1)/2-s} e^{-t} \int_W |x|^{2s-(n-1)} f(x) dx.$$

This proves the functional equation. □

Theorem 10 is now an immediate consequence of equation (6) and of the functional equation in Lemma 12 (applied to the value $s = \frac{(n-1)(1-u)}{2}$).

Notation 1. We denote by $\widehat{\pi_u}$ the **completion** of the space $\mathcal{S}(W)$ under the metric defined by Theorem 10. We then get a metric on $\widehat{\pi_u}$. The G -invariance of the pairing between $\mathcal{F}_u$ and $\mathcal{F}_{-u}$ and the definition of the inner product in Theorem 10 implies that the metric on $\widehat{\pi_u}$ is G -invariant. Note that elements of the completion $\widehat{\pi_u}$ may

not be measurable functions on G , but only “generalised functions” or distributions.

3.2. Embedding of the unramified Complementary series.

Notation 2. Consider $H = O(n-1, 1) \subset O(n, 1) = G$ embedded by fixing the $n-1$ -th basis vector e_{n-1} . Denote by $\sigma_{u'}$ the representation $\pi_{u'}$ constructed in the last section, not for the group G , but, for the group H . We set $u' = \frac{(n-1)u-1}{n-2}$. We assume that $0 < u' < 1$ which means that $\frac{1}{n-1} < u < 1$.

Denote by W the space $\mathbb{R}^{n-1}$. Let $W' \subset W$ be the subspace $\mathbb{R}^{n-2}$ whose last co-ordinate in $\mathbb{R}^{n-1}$ is zero.

Let $\widehat{W}_u$ be the completion of the Schwarz space of W under the metric

$$|\phi|_u^2 = \int_W \frac{dx}{|x|^{(n-1)u}} |\phi(x)|^2.$$

The convergence of this integral near infinity is clear since ϕ lies in the Schwartz space; the convergence of the integral near 0 follows by converting to polar co-ordinates and by using the assumption that $0 < u < 1$.

Define the space $\widehat{W}'_{u'}$ correspondingly for W' as the completion of the Schwartz space of W' under the metric

$$|\psi|_{u'}^2 = \int_{W'} \frac{dy}{|y|^{(n-2)u'}} |\psi(y)|^2.$$

Now, $\widehat{W}_u$ consists of all measurable functions ϕ on $W = \mathbb{R}^{n-1}$ which the u -norm, that is, integral

$$|\phi|_u^2 = \int_W \frac{dx}{(|x|)^{(n-1)u}} |\phi(x)|^2,$$

is finite. One may define similarly, the u' -norm of a function on W' . Given ϕ in the space $\mathcal{S}(W)$, we define the function $J(\phi)$ on W by setting. for all $(y, t) \in W' \times \mathbb{R} = \mathbb{R}^{n-2} \times \mathbb{R} \simeq \mathbb{R}^{n-1} = W$,

$$J(\phi(y, t)) = \phi(y).$$

Then $J(\phi)$ is a bounded measurable function on $\mathbb{R}^{n-1}$.

Proposition 13. *The map J defined on the Schwartz space $\mathcal{S}(W')$ has its image in the Hilbert space $\widehat{W}_u$ and extends to an isometry from $\widehat{W}'_{u'} \rightarrow \widehat{W}_u$.*

Proof. We compute the u -norm of the bounded function $J(\phi)$:

$$\|J(\phi)\|_u^2 = \int_{W' \times \mathbb{R}} \frac{dy \, dt}{(|y|^2 + t^2)^{\frac{(n-1)u}{2}}} |\phi(y)|^2.$$

By changing the variable t to $|y|t$ and using the definition of the map J , we see that this u -norm is equal to the integral

$$\int \frac{dt}{(1+t^2)^{\frac{(n-1)u}{2}}} \int_{W'} \frac{dy}{|y|^{(n-1)u-1}} |\phi(y)|^2.$$

The integral over t converges when the exponent of the denominator term $\frac{(n-1)u}{2}$ is strictly larger than 1; that is, $u > \frac{1}{n-1}$. The integral over y is simply the u' norm of ϕ since $(n-1)u-1 = (n-2)u'$. Therefore, the u -norm of $J(\phi)$ is the u' -norm of ϕ up to a constant (the integral over the variable t of the function $\frac{1}{(1+t^2)^{(n-1)u}}$). We have therefore proved the proposition. $\square$

The completion $\widehat{\pi}_u$ (and similarly $\widehat{\sigma}_{u'}$) was defined with respect to the metric $\langle \phi, I(u)(\phi) \rangle$ (we use Theorem 10 to say that this is indeed a metric). It is clear from this definition that for $\phi \in \mathcal{S}(W) \subset \mathcal{F}_u$, the map $\phi \mapsto \widehat{\phi}$ (the roof over ϕ refers to the Fourier transform) gives an isometry from $\widehat{\pi}_u$ onto $\widehat{W}_u$. Similarly $\widehat{\sigma}_{u'}$ is isometric to $\widehat{W}'_{u'}$. Therefore, Proposition 13 says that the Hilbert space $\widehat{\sigma}_{u'}$ is isometrically embedded in the Hilbert space $\widehat{\pi}_u$. We now show that the map J is equivariant with respect to the action of the subgroup H on both sides.

Notation 3. Let $\phi^* \in \mathcal{S}(W) \subset \mathcal{F}_u$ be a function whose Fourier transform is a smooth compactly supported function $\widehat{\phi}^*$ on $\mathbb{R}^{n-1} \setminus \{0\}$. Denote by $\phi \in \mathcal{F}_{-u}$ the image of ϕ^* under the map $I(u)$. By the definition of $I(u)$, we have the formula (see Lemma 11),

$$\phi(u(y)w) = \int_{\mathbb{R}^{n-1}} \phi^*(u(y)wu(x)w) dx.$$

Using Lemma 12, we now get

$$\phi(y) = \int_{\mathbb{R}^{n-1}} \frac{dx}{|x|^{(n-1)u}} \widehat{\phi}^*(x) e^{-2ix \cdot y}.$$

Denote by $\mathcal{A}_G$ the map $\widehat{\phi}^* \mapsto \phi$ from the space $\mathcal{C}_c^\infty(\mathbb{R}^{n-1} \setminus \{0\})$ into the space $\mathcal{F}_{-u}$, by the preceding formula. By our identifications, this extends to an isometry $\mathcal{A}_G$ from $\widehat{W}_u$ onto the Hilbert space $\widehat{\pi}_{-u} \simeq \widehat{\pi}_u$. This map exists for $0 < u < 1$. The G action on the image $\widehat{\pi}_{-u}$ gives, via this isomorphism $\mathcal{A}_G$, the G -action on the Hilbert space $\widehat{W}_u$.

We similarly have a map $\mathcal{A}_H$ from the space $\mathcal{C}_c^\infty(\mathbb{R}^{n-2} \setminus \{0\})$ into $\mathcal{F}_{-u'}^H$ (of corresponding functions on the group H) which extends to an isometry from the space $\widehat{W}'_{-u'}$ onto $\widehat{\sigma}_{-u'} \simeq \widehat{\sigma}_{u'}$. The H action on $\widehat{\sigma}_{-u'}$ gives an action of H on the space $\widehat{W}'_{-u'}$.

Let $J^* : \widehat{W}_u \rightarrow \widehat{W}_{u'}$ be the adjoint of the map J .

Proposition 14. *We have the formula for the adjoint J^* of J :*

$$J^* = \mathcal{A}_H^{-1} \circ \text{res} \circ \mathcal{A}_G.$$

(Here $\text{res} : \mathcal{F}_{-u} \rightarrow \mathcal{F}_{-u'}^H$ is simply restricting the functions on G to the subgroup H (since $(n-1)u - 1 = (n-2)u'$, it follows that the restriction maps the functions in $\mathcal{F}_{-u}$ into the functions in $\mathcal{F}_{-u'}^H$)).

Proof. The adjoint is defined by the formula

$$\langle g, J^* f \rangle_{W_{u'}} = \langle Jg, f \rangle_{W_u},$$

for all functions $f \in \mathcal{C}_c^\infty(\mathbb{R}^{n-1} \setminus \{0\})$ and $g \in \mathcal{C}_c^\infty(\mathbb{R}^{n-2} \setminus \{0\})$. We compute the right hand side of this formula. By definition, the inner product of Jg and f is the integral

$$\int_{\mathbb{R}^{n-1}} \frac{dx}{|x|^{(n-1)u}} Jg(x) f(x).$$

Write $x \in \mathbb{R}^{n-1} = \mathbb{R}^{n-2} \times \mathbb{R}$ as $x = (y, s)$. Then, by the definition of J , $Jg(x) = Jg(y, s) = g(y)$ and the preceding integral is the integral

$$\int_{\mathbb{R}^{n-2}} dy g(y) \int ds f(y, s) \frac{1}{(|y|^2 + s^2)^{(n-1)u}}.$$

By changing s to $|y|s$, the integral becomes

$$\int_{\mathbb{R}^{n-2}} \frac{dy}{|y|^{(n-1)u-1}} g(y) \int_{\mathbb{R}} \frac{ds}{(1+s^2)^{(n-1)u/2}} f(y, |y|s).$$

By the definition of the $\widehat{W}'_{u'}$ inner product, this is $\langle g, h \rangle_{W'_{u'}}$ where h is the function in y defined by

$$h(y) = \int_{\mathbb{R}} \frac{ds}{(1+s^2)^{(n-1)u/2}} f(y, |y|s).$$

which is the integral over the variable s in the preceding equality. We have proved that $\langle Jg, f \rangle = \langle g, h \rangle$.

By the definition of the adjoint, this means that $h = J^* f$. We have therefore

$$J^*(f) = \int_{\mathbb{R}} \frac{ds}{(1+s^2)^{(n-1)u/2}} f(y, |y|s).$$

We now compute $\mathcal{A}_H J^*(f)(x)$ at a point $x \in \mathbb{R}^{n-2}$. By the definition of $\mathcal{A}_H$, this is the integral

$$\int_{\mathbb{R}^{n-2}} \frac{dy}{|y|^{(n-2)u'}} J^*(f)(y) e^{-2ix \cdot y}$$

The above integral formula for J^* shows that $\mathcal{A}_H J^*(f)(x)$ is the integral

$$\mathcal{A}_H J^* f(x) = \int_{\mathbb{R}^{n-2} \times \mathbb{R}} \frac{dy \, ds}{(|y|)^{(n-2)u'} (1+s^2)^{(n-1)u/2}} f(y, |y|s) e^{-2ix \cdot y}.$$

Changing the variable s back to $\frac{s}{|y|}$ and substituting it in this integral (and using the fact that $(n-2)u' = (n-1)u - 1$), we get

$$\mathcal{A}_H J^* f(x) = \int_{\mathbb{R}^{n-2} \times \mathbb{R}} \frac{dy \, ds}{(|y|^2 + s^2)^{(n-1)u/2}} f(y, s) e^{-2i(x,0) \cdot (y,s)}.$$

The latter is simply $\mathcal{A}_G(f)(x)$ for $x \in \mathbb{R}^{n-2}$. By taking f to be $\mathcal{A}_G^{-1}\phi$, we get, for all $x \in \mathbb{R}^{n-2}$, and all ϕ which are $\mathcal{A}_G$ -images of compactly supported smooth function on $\mathbb{R}^{n-1} \setminus \{0\}$, that

$$\mathcal{A}_H J^* \mathcal{A}_G^{-1}(\phi)(x) = \phi(x) = \text{res}(\phi)(x).$$

This proves the Proposition. $\square$

Corollary 1. *The isometric map $J : \widehat{W}'_{u'} \rightarrow \widehat{W}_u$ is equivariant for the action of H on both sides.*

Proof. It is enough to prove that the adjoint J^* of J is equivariant for H action. It follows from the Proposition, that $J^* = \mathcal{A}_H^{-1} \circ \text{res} \circ \mathcal{A}_G$. All the maps $\mathcal{A}_H$, $\mathcal{A}_G$ and res are H -equivariant. The Corollary follows. $\square$

Corollary 2. *The map $\pi_{-u} \rightarrow \sigma_{-u'}$ from the module of K -finite vectors in $\widehat{\pi}_{-u}$ onto the corresponding vectors in $\widehat{\sigma}_{-u'}$ is continuous.*

Proof. The map res is, by the Proposition, essentially the adjoint J^* of the map J . But J is an isometry which means that J^* is a continuous projection. Therefore, res is continuous. $\square$

Theorem 15. *Let $\frac{1}{n-1} < u < 1$ and $u' = \frac{(n-1)u-1}{n-2}$. The complementary series representation $\widehat{\sigma}_{u'}$ of $SO(n-1, 1)$ occurs discretely in the restriction of the complementary series representation $\widehat{\pi}_u$ of $SO(n, 1)$.*

Proof. We have the H -equivariant isometry from $\widehat{W}'_u$ into $\widehat{W}_u$. The former space is by construction, isometric to $\sigma_{u'}$ and the latter, to $\widehat{\pi}_u$.

Therefore, the Theorem follows. $\square$

4. RAMIFIED COMPLEMENTARY SERIES

In this section, we obtain an embedding of the ramified complementary series representation of $H = SO(n-1, 1)$ in the ramified complementary series representation of $G = SO(n-1, 1)$. The method follows that for the unramified case, except that in the unramified case, the G -invariant pairing on the complementary series was directly proved to be positive definite, using Theorem 10. However, in the ramified case, this seems more difficult to prove. But, the positive definiteness is actually known by [Ba-B]; given this positive definiteness, the proof proceeds as in the unramified case. Nevertheless, we give complete details, since we will have to deal not with scalar valued operators, but with matrix valued ones, and some complications arise.

4.1. A Model for Ramified Complementary Series. Let $1 \leq i \leq [n/2]$ be an integer. Set $W = \mathbb{R}^{n-1}$. Denote by $\mathcal{S}(W, i)$ the space of functions on W of Schwartz class, with values in the vector space $\wedge^i = \wedge^i \mathfrak{p}_M$ (recall that $\mathfrak{p}_M \simeq \mathbb{C}^{n-1}$ is the standard representation of $M = SO(n-1)$ twisted by a character on $O(n-1)$).

As we saw before, $G/P = K/M$ and the map $x \mapsto u(x)wP \in G/P$ maps W injectively onto the “big Bruhat Cell” in G/P . Let $u \in \mathbb{C}$. Denote by $\mathcal{F}_u(i)$ the space of **continuous** functions on G with values in $\wedge^i$ which transform according to the formula

$$f(gman) = \wedge^i(m)(f(g))\rho_P(a)^{-(1+u)},$$

for all $(g, m, a, n) \in G \times A \times M \times N$.

An element in the Schwartz space $\mathcal{S}(W, i)$ may be extended to a section of the vector bundle on G/P associated to $\wedge^i$ (that is, an element of $\mathcal{F}_u(i)$). We denote this embedding by $E(u) : \mathcal{S}(W, i) \subset \mathcal{F}_u(i)$.

Lemma 16. *If $\operatorname{Re}(u) > 0$ and $f \in \mathcal{F}_u(i)$, then the integral*

$$I_i(u)(f)(g) = \int_W dx f(gu(x)w),$$

converges and as a function of $g \in G$, lies in $\mathcal{F}_{-u}(i)$.

Proof. The boundedness of M shows that the convergence amounts to the convergence (for $\operatorname{Re}(u) = C$) of the integral

$$\int_W \frac{dx}{(1 + |x|^2/2)^{(n-1)(1+C)/2}};$$

the latter integral converges because of the assumption $C > 0$. □

Given $f \in \mathcal{S}(W, i) \subset \mathcal{F}_u(i)$, we can then form the inner product (in the vector space $\wedge^i$) of the vectors $f((u(y)w)$ and $I_i(u)(f)(u(y)w)$. By an abuse of notation, we write $f(y) = f(u(y)w)$. Then the inner product is the function $x \mapsto \langle f(y), I_i(u)(f)(y) \rangle$. Since $f(y)$ is in the Schwartz space and $I_i(u)f(y)$ is bounded on W , it follows that this inner product function is integrable over y and thus we get a pairing

$$\langle f, I_i(u)f \rangle = \int_W dy \langle f(y), I_i(u)(f)(y) \rangle.$$

This is a G -invariant bilinear form on $\mathcal{S}(W, i)$. We now use a Theorem of [Ba-B] to obtain that if $0 < u < \frac{2i}{n-1}$, then this pairing is a positive definite inner product on $\mathcal{S}(W, i)$.

Recall from Lemma 4 that if $x \in \mathbb{R}^{n-1}$, then the Bruhat decomposition of an element of the form

$$wu(x)w = u\left(\frac{2x}{|x|^2}\right)wm(x)d(a)u(x),$$

with $m(x) = 1 - \frac{2^t xx}{|x|^2} \in M$ being the reflection about the orthogonal complement of x , and $a = \frac{|x|^2}{2}$ defining the diagonal matrix $d(a)$.

We now prove a Lemma analogous to Lemma 11.

Lemma 17. *If $\operatorname{Re}(u) > 0$ then the formula for the inner product is given, for $f \in \mathcal{F}_u$ and $y \in \mathbb{R}^{n-1}$, by the formula*

$$(I(u)f)(wu(y)) = \int_{\mathbb{R}^{n-1}} \frac{dx}{|x|^{(n-1)(1-u)}} m(x)(f(y+x)).$$

In this equation, the function f has values in the representation space $\wedge^i$ of the group M and $m(x)$, being an element of M , acts on the vector $f(u(y+x)w)$, which, by an abuse of notation, is written $f(y+x)$.

Proof. The intertwining operator is

$$I_i(u)f(u(y)w) = \int_{\mathbb{R}^{n-1}} \frac{dx}{|x|^{(n-1)(1-u)}} f(u(y)wu(x)w).$$

We now use Lemma 4 to conclude that

$$I_i(u)f(y) = \int_{\mathbb{R}^{n-1}} dx \wedge^i(m(x))(f(y + \frac{2x}{|x|^2})).$$

The Lemma now follows from Lemma 5, since

$$m\left(\frac{2x}{|x|^2}\right) = m(x) :$$

the two reflections are the same since the orthogonal complements are the same. $\square$

In the following we denote by $m_i(x)$ the linear operator $\wedge^i(m(x))$ acting on the representation space $\wedge^i$ of M .

We now use Lemma 8 to conclude the following. Given $f \in \mathcal{F}_u$, the pairing of $\mathcal{F}_u(i)$ and $\mathcal{F}_{-u}(i)$, is given by

$$L(< f(y), I_i(u)f(y) >_{\wedge^i}) = \int_{\mathbb{R}^{n-1}} dy < f(y), I_i(u)f(y) >_{\wedge^i}.$$

Using the formula for the intertwining operator in Lemma 17, we then have

Lemma 18. *The inner product of f with itself in $\mathcal{F}_u(i)$ is given by*

$$< f, f >_{\mathcal{F}_u(i)} = \int_{\mathbb{R}^{n-1}} dy < f(y), \int_{\mathbb{R}^{n-1}} \frac{dx}{|x|^{(n-1)(1-u)}} m_i(x)(f(y+x)) >.$$

If we now assume that $f \in \mathcal{S}(W, i) \subset \mathcal{F}_u(i)$, then we see that the double integral in the foregoing Lemma can be interchanged. The integral of a product of Schwartz class functions is the integral of their Fourier transforms. The Fourier transform of a translate of a function by a vector is the multiple of the transform of the function by a character. Using these observations (the Fourier transform of $f(y)$ is denoted $\widehat{f}(y)$), we have, for $\phi \in \mathcal{S}(W, i)$, the formula

$$\begin{aligned} < \phi, I_i(u)\phi >_{\mathcal{S}(W, i)} = \\ = \int_{\mathbb{R}^{n-1}} \frac{dx}{|x|^{(n-1)(1-u)}} \int dy < \widehat{\phi}(y), m_i(x)(\widehat{\phi}(y)) > e^{-2iy \cdot x}. \end{aligned}$$

Definition 1. Set $W = \mathbb{R}^{n-1}$ and $\mathcal{S}(W, i)$ the space of $\wedge^i$ valued Schwartz class functions on W , as before. We denote by $< \phi, \phi >_{W_u}$, for $\phi \in \mathcal{S}(W, i)$, the sesquilinear form

$$< \phi, \phi >_{W_u(i)} = \int_W \frac{dx}{|x|^{(n-1)(1-u)}} \int_W dy < \phi(y), m_i(x)(\phi(y)) >_{\wedge^i} e^{-2iy \cdot x}.$$

Since the map $\phi \mapsto E(u)(\widehat{\phi}) \in \mathcal{F}_u(i)$ defined in section (4.1) is an injection, it follows from the result of [Ba-B] quoted earlier that the above sesquilinear form is positive definite, provided u is real and $0 < u < 1 - \frac{2i}{n-1}$. Denote by $\widehat{W}_u(i)$ the **completion** of the Schwartz space $\mathcal{S}(W, i)$ with respect to this inner product. We have then an isometry $\widehat{\pi}_u(i) \rightarrow \widehat{W}_u(i)$, which, on Schwartz class functions $\phi \simeq E(u)(\phi)$ in $\widehat{\pi}_u(i)$, is given by the Fourier transform $\phi \mapsto \widehat{\phi}$.

Definition 2. We define analogously, the subspace $W' = \mathbb{R}^{n-2}$ and if $0 < u' < 1 - \frac{2i}{n-2}$, we get a metric on the Schwartz space $\mathcal{S}(W', i)$ given by

$$\langle \phi, \phi \rangle_{W', (i)} = \int_{W'} \frac{dx}{|x|^{(n-2)(1-u')}} \int_{W'} dy \langle \phi(y), m_i(x)(\phi(y)) \rangle e^{-2ix \cdot y}.$$

The completion is denoted $\widehat{W}'_{u'}(i)$. This is isometric to the representation $\widehat{\sigma}_{u'}(i)$ of $H = SO(n-2, 1)$.

4.2. Embedding of the Ramified Complementary Series.

Notation 4. We have the subgroup $H = SO(n-1, 1) \subset G = SO(n, 1)$; the embedding is so that the intersection $M_H = M \cap H$ is the compact part of the Levi subgroup of the minimal parabolic subgroup $P \cap H$ of H ; we have the representation $\mathfrak{p}_{M_H}$ and we have a natural inclusion $\mathfrak{p}_{M \cap H} \subset \mathfrak{p}_M$ induced by the inclusion of the Lie algebra $\mathfrak{h}$ of H in the Lie algebra $\mathfrak{g}$ of G . Consequently, we have the inclusion $\wedge^i \mathfrak{p}_{M_H} \subset \wedge^i \mathfrak{p}_M$ and functions with values in the subspace $\wedge^i \mathfrak{p}_{M_H}$ may be viewed as functions with values in the space $\wedge^i \mathfrak{p}_M$.

We define a map $J : \widehat{W}'_{u'}(i)$ into $\wedge^i$ valued continuous functions on $W = \mathbb{R}^{n-1} = W' \times \mathbb{R}$, by the formula

$$J(\phi)(y, t) = \phi(y),$$

for all $(y, t) \in \mathbb{R}^{n-2} \times \mathbb{R} = W' \times \mathbb{R} = W$. Using the convention of the preceding paragraph, we have viewed $\phi : W' \rightarrow \wedge^i \mathfrak{p}_{M \cap H}$ as taking values in the larger vector space $\wedge^i \mathfrak{p}_M$.

Lemma 19. *Suppose that $\frac{1}{n-1} < u < 1 - \frac{2i}{n-1}$ and that $(n-2)(1-u') = (n-1)(1-u)$. The map J extends to an isometry from $\widehat{W}'_{u'}(i)$ into the Hilbert space $\widehat{W}_u(i)$.*

Proof. It is enough to check that for ϕ in the Schwartz space $\mathcal{W}'_{u'}(i)$, the function $J(\phi)$ has finite $\widehat{W}_u(i)$ norm and that its norm is the norm of ϕ as an element of $\widehat{W}'_{u'}(i)$.

We then compute the latter norm. Set $\lambda = (n-1)(1-u)$. By assumption, $\lambda = (n-2)(1-u')$. Recall that $m_i(x, t)$ is the i -th exterior power of the element $m(x, t) \in M$ and is viewed as an endomorphism of $\wedge^i \mathfrak{p}_M$. We have

$$\begin{aligned} \langle J\phi, J\phi \rangle_{W_u(i)} &= \int_{W' \times \mathbb{R}} \frac{dx \, dt}{(\sqrt{|x|^2 + t^2})^\lambda} \\ &\int_{W' \times \mathbb{R}} dy \, ds \langle \phi(y), m_i(x, t)\phi(y) \rangle e^{-2ix \cdot y} e^{-2is \cdot t}. \end{aligned}$$

We have used here the fact that $J\phi(y, s) = \phi(y)$. The assumption that ϕ is of Schwartz class implies that all these integrals can be interchanged, and hence we may write

$$\langle J\phi, J\phi \rangle = \int_{W' \times W'} \frac{dx \, dy}{|x|^\lambda} \langle \phi(y), M_i(x)(\phi(y)) \rangle_{\wedge^i} e^{-2ix \cdot y},$$

where $M_i(x)$ is the linear operator on the finite dimensional vector space $\wedge^i$ given by the integral

$$M_i(x) = \int ds (\mathcal{F}_t \frac{m_i(x, t)}{(\sqrt{|x|^2 + t^2})^\lambda})(s),$$

where $\mathcal{F}_t$ denotes the Fourier transform with respect to the real variable t .

Recall that $m_i(x, t)$ is the i -th exterior power of the matrix $1 - 2\frac{taa}{|a|^2}$ where $a = (x, t)$ is the row matrix of size $n - 1$. Therefore, the product $\Phi_i(x, t) = (|x|^2 + t^2)^i m_i(x, t)$ is a homogeneous polynomial of degree $2i$ on the entries of the row matrix (x, t) .

We may write

$$(7) \quad M_i(x) \Gamma(\lambda/2 + i) = \int_0^\infty \frac{dv}{v} v^{\lambda/2 + i} e^{-v} \int_{\mathbb{R}} ds \left(\int_{\mathbb{R}} \frac{dt}{(\sqrt{|x|^2 + t^2})^\lambda} \Phi_i(x, t) e^{-2ist} \right).$$

Changing v to $v(|x|^2 + t^2)$, we obtain

$$M_i(x) = \frac{1}{\Gamma(\lambda/2 + i)} \int_0^\infty \frac{dv}{v} v^{\lambda/2 + i} e^{-v|x|^2} \int_{\mathbb{R}} ds \left(\int_{\mathbb{R}} dt \Phi_i(x, t) e^{-vt^2} e^{-2ist} \right).$$

Set

$$G_i(s, v) = \int dt e^{-vt^2} \Phi_i(x, t) e^{-2\sqrt{-1}st}.$$

Changing t to $\frac{t}{\sqrt{v}}$ in this integral, we obtain that

$$G_i(s, v) = \frac{1}{\sqrt{v}} \int_{\mathbb{R}} dt e^{-t^2} e^{-2\sqrt{-1}st/\sqrt{v}} \Phi_i(x, t/\sqrt{v}).$$

Thus $G_i(v, s)$ is essentially the Fourier transform of some function evaluated at $s/\sqrt{v}$.

The Fourier transform of e^{-t^2} is itself and the Fourier transform of the product of e^{-t^2} with a power t^j of t is the j -th derivative of e^{-t^2} . The j th derivative is of the form $e^{-t^2} P_j(t)$ where P_j is a polynomial of degree j . Since $\Phi_i(x, t)$ is a homogeneous polynomial of degree $2i$, it follows that for some polynomial P_i^* of degree i , we have

$$G_i(s, v) = \frac{1}{\sqrt{v}} \frac{1}{v^i} P_i^*(s/\sqrt{v}) e^{-s^2/v}.$$

Then we get

$$\begin{aligned}
& \int_0^\infty \frac{dv}{v} v^{\lambda/2+i} e^{-v|x|^2} \int ds G_i(v, s) = \\
& \text{(after changing } s \text{ to } s\sqrt{v}) \\
& = \int_0^\infty \frac{dv}{v} e^{-|x|^2} v^{\lambda/2} \int_{\mathbb{R}} ds P_i^*(s) e^{-s^2} = \\
& = \frac{1}{|x|^\lambda} \Gamma(\lambda/2) C,
\end{aligned}$$

where C is some bounded number.

Therefore, in the triple integral defining $M_i(x)$ in the equation (7), we see that the integrand is an integrable function in t, s, v ; therefore its integral over t is an **integrable function** in s and v ; the integral over t gives an integrable function over s whose integral is (by Fourier inversion), the value of the function at $t = 0$. We therefore get

$$M_i(x) = \frac{1}{\Gamma(\lambda/2 + i)} \int_0^\infty \frac{dv}{v} v^{\lambda/2+i} e^{-v|x|^2} |x|^{2i} m_i(x, 0) = \frac{m_i(x, 0)}{|x|^\lambda},$$

(the last equality follows by changing v to $v/(|x|)^2$).

Plugging in this value for $M_i(x)$ in the expression for $\langle J\phi, J\phi \rangle$ above, we obtain

$$\langle J\phi, J\phi \rangle = \int_{W'} \frac{dx}{|x|^\lambda} \int_{W'} dy \langle \phi(y), m_i(x, 0) \phi(y) \rangle e^{-2ix \cdot y}.$$

By the definition of norm on $\widehat{W}'_{u'}(i)$ this is $\langle \phi, \phi \rangle_{W'_{u'}(i)}$, since **by assumption**, $\lambda = (n-1)(1-u) = (n-2)(1-u')$.

Therefore, $J\phi$ has finite norm and its norm is equal to the norm of ϕ . This proves the Lemma. $\square$

Notation 5. Denote by $\mathcal{A}_G = \mathcal{A}_G(u)$ the composite of the two maps: $\phi \mapsto \widehat{\phi}$ from $\mathcal{S}(W, i)$ into $E(u)(\mathcal{S}(W, i) \subset \widehat{\pi}_u(i))$ and the intertwining map $I_i(u) : E(u)(\mathcal{S}(W, i)) \rightarrow \mathcal{F}_{-u}(i)$. If $0 < u < 1 - \frac{2i}{n-1}$, then $I_i(u)$ is an isomorphism, and we get a metric on the image of $I_i(u)$ whose completion is denoted $\widehat{\pi}_{-u}(i)$; the map $I_i(u)$ yields an isometry $I_i(u) : \widehat{\pi}_u(i) \rightarrow \widehat{\pi}_{-u}(i)$.

The composite of the Fourier transform with this isometry $I_i(u)$ gives an isometry

$$\mathcal{A}_G = \mathcal{A}_G(u) : \widehat{W}_u(i) \rightarrow \widehat{\pi}_{-u}(i).$$

The G -action on $\widehat{W}_u(i)$ is borrowed from $\widehat{\pi}_u(i)$ via the isometry induced by the Fourier transform. Therefore, the map $\mathcal{A}_G$ is a G -equivariant

map.

We have similarly the isometry

$$\mathcal{A}_H = \mathcal{A}_H(u') : \widehat{W}'_{u'}(i) \rightarrow \widehat{\sigma}_{-u'}(i),$$

Similarly, the map $\mathcal{A}_H$ is an H -equivariant isometry.

Notation 6. Given that J is an isometry from $\widehat{W}'_{u'}(i)$ into $\widehat{W}_u(i)$, denote by J^* its adjoint; J^* is the orthogonal projection from $\widehat{W}_u(i)$ onto $\widehat{W}'_{u'}(i)$.

The functions in $\mathcal{F}_{-u}(i)$ have values in the representation $\wedge^i \mathfrak{p}_M$ of the group $M = SO(n-1)$ and have the tranformation property

$$f(gmd(a)n) = \wedge^i(m)(f(g))\rho_P(d(a))^{-(1-u)}.$$

Here, if $d(a)$ is the diagonal matrix in $SO(n, 1)$ corresponding to $a \in \mathbb{R}^*$ with $a > 0$, then $\rho_P(d(a)) = a^{n-1}$. Consequently,

$$\rho_P(d(a))^{-(1-u)} = a^{(n-1)(1-u)} = a^{(n-2)(1-u')}.$$

Denote by $\wedge^i \mathfrak{p}_{M_H}$ the representation of $M \cap H = SO(n-2)$ on the i -th exterior power of its representation on $\mathfrak{p}_{M \cap H}$, and view $\wedge^i \mathfrak{p}_{M_H}$ as an $M \cap H$ -equivariant *quotient* of the representation $\wedge^i \mathfrak{p}_M$ of M . There is no conflict with the previous notation where $\wedge^i \mathfrak{p}_{M \cap H}$ was viewed as a *subspace* of $\wedge^i \mathfrak{p}_M$, since it occurs with multiplicity one in the representation $\wedge^i \mathfrak{p}_M$, of $M \cap H$.

Consequently, we have a restriction map (actually, restriction followed by the above quotient map)

$$res : \mathcal{F}_{-u}(i) \rightarrow \mathcal{F}_{-u'}(i),$$

which is H -equivariant.

We now prove an analogue of Proposition 14, for the *ramified* case.

Proposition 20. *Suppose that $\frac{1}{n-1} < u < 1 - \frac{2i}{n-1}$ and define u' by $(n-1)(1-u) = (n-2)(1-u')$. We have the equality*

$$\mathcal{A}_H \circ J^* = res \circ \mathcal{A}_G,$$

with $\mathcal{A}_G = \mathcal{A}_G(u)$ and $\mathcal{A}_H = \mathcal{A}_H(u')$.

Proof. Let $\phi, \psi \in \mathcal{S}(W, i)$. Since $\mathcal{A}_G$ is an isometry, we have the equality

$$\langle \phi, \psi \rangle_{\widehat{W}_u(i)} = \langle \mathcal{A}_G(\phi), \mathcal{A}_G(\psi) \rangle_{\mathcal{F}_{-u}(i)}.$$

Now $\mathcal{A}_G(\phi) = I(u) \circ \widehat{\phi}$. If $\langle, \rangle_{\mathcal{F}_u \times \mathcal{F}_{-u}}$ denotes the pairing between $\mathcal{F}_u(i)$ and $\mathcal{F}_{-u}(i)$, then we get

$$\langle \phi, \psi \rangle_{\widehat{W}_u(i)} = \langle \widehat{\phi}, \mathcal{A}_G \psi \rangle_{\mathcal{F}_u \times \mathcal{F}_{-u}}.$$

By Lemma 8, this is equal to

$$\int_W dx \langle \phi(x), (\mathcal{A}_G \psi)(x) \rangle_{\wedge^i},$$

which is the L^2 inner product of the functions involved. We have therefore proved

$$(8) \quad \langle \phi, \psi \rangle_{\widehat{W}_u(i)} = \langle \widehat{\phi}, (\mathcal{A}_G \psi) \rangle_{L^2(W)}.$$

Similarly, we have for the subgroup H and $\phi, \psi \in \widehat{W}'_{u'}(i)$, the formula

$$(9) \quad \langle \phi, \psi \rangle_{\widehat{W}'_{u'}(i)} = \langle \widehat{\phi}_1, (\mathcal{A}_G \psi) \rangle_{L^2(W')},$$

where $\widehat{\phi}_1$ is the Fourier transform for the subspace W' . From equation(8) we get for $\phi \in \mathcal{W}'(i)$,

$$\begin{aligned} \langle J\phi, \psi \rangle_{W_u(i)} &= \langle \widehat{J\phi}, \mathcal{A}_G \psi \rangle_{L^2(W)} = \langle \phi, \widehat{\mathcal{A}_G \psi} \rangle_{L^2(W)} = \\ &= \int_{W' \times \mathbb{R}} dx dt \langle \phi(x), (\widehat{\mathcal{A}_G \psi})(x, t) \rangle. \end{aligned}$$

The Fourier transform on $W' \times \mathbb{R}$ is the Fourier transform on W' followed by the Fourier transform in the variable $t \in \mathbb{R}$. Moreover, the integral of the Fourier transform in t is the value of the function at 0 (Fourier inversion). This shows that the last integral is equal to

$$\langle J\phi, \psi \rangle = \int_{W'} dx \langle \phi(x), \widehat{\mathcal{A}_G \psi}_1(x, 0) \rangle = \langle \widehat{\phi}_1, \text{res} \circ \mathcal{A}_G \psi \rangle_{L^2(W')}.$$

On the other hand, from the definition of the adjoint J^* and equation (9), we get

$$\langle J\phi, \psi \rangle_{W_u(i)} = \langle \phi, J^* \psi \rangle_{W'_{u'}(i)} = \langle \widehat{\phi}_1, \mathcal{A}_H \circ J^* \psi \rangle_{L^2(W')}.$$

The last two equations show that

$$\mathcal{A}_H \circ J^* \psi = \text{res} \circ \mathcal{A}_G \psi,$$

for all Schwartz functions ψ on W . Therefore, the Proposition follows. $\square$

Corollary 3. *The map $J : \widehat{W}'_{u'}(i) \rightarrow \widehat{W}_u(i)$ is equivariant for the action of H on both sides.*

Proof. By the Proposition 20, the map J^* satisfies $\mathcal{A}_H \circ J^* = res \circ \mathcal{A}_G$. All the maps $\mathcal{A}_H$, $\mathcal{A}_G$ and res are equivariant for H . The corollary follows. $\square$

Corollary 4. *The restriction map $res : \mathcal{S}(W_{-u}(i)) \rightarrow \mathcal{S}(W'_{-u'}(i))$ is continuous for the G and H invariant metrics on the Schwartz spaces $\mathcal{S}(W_{-u}, i)$ and $\mathcal{S}(W'_{-u'}, i)$.*

Proof. The map $res = \mathcal{A}_H \circ J^* \circ \mathcal{A}_G^{-1}$; the maps $\mathcal{A}_G, \mathcal{A}_H$ are isometries and J^* is an orthogonal projection (adjoint of an isometry J). Consequently, res is also an orthogonal projection and is in particular, continuous. $\square$

Theorem 21. *The representation on the Hilbert space $\widehat{\sigma_{u'}(i)}$ occurs discretely in the Hilbert space $\widehat{\pi_u(i)}$ if $\frac{1}{n-1} < u < 1 - \frac{2i}{n-1}$.*

Proof. We have in fact (Corollary 4) that the map

$$res : \pi_{-u} \rightarrow \sigma_{-u'}$$

is continuous for the norms on the representations involved. $\square$

5. THE MAIN THEOREM AND COROLLARIES

Theorem 22. *Let $n \geq 3$, $H = SO(n-1, 1)$ and $G = SO(n, 1)$. Let i be an integer with $0 \leq i \leq [n/2] - 1$ where $[x]$ denotes the integral part of x . Let $\frac{1}{n-1} < u < 1 - \frac{2i}{n-1}$. Then the complementary series representation*

$$\widehat{\pi}_u(i) = \text{Ind}_P^G(\wedge^i \mathfrak{p}_M \otimes \rho_P(a)^u),$$

contains discretely, the complementary series representation

$$\widehat{\sigma}_{u'}(i) = \text{Ind}_{P \cap H}^H(\wedge^i \mathfrak{p}_{M \cap H} \otimes \rho_{P \cap H}^{u'}),$$

where $u' = \frac{(n-1)u-1}{n-2}$. Further, as u tends to $1 - \frac{2i}{n-1}$, u' tends to $1 - \frac{2i}{n-2}$.

Proof. We have already proved this (see Theorem 21) except for the last statement, which is trivial to prove. $\square$

Corollary 5. *Let $i \leq [n/2] - 1$. Then the cohomological representation $A_i(n)$ of $SO(n, 1)$ of degree i restricted to $SO(n-1, 1)$ contains discretely, the cohomological representation $A_i(n-1)$.*

The discrete series representation $A_i(2i)$ of $SO(2i, 1)$ is contained discretely in the representation A_i .

Proof. To prove the first part, we use Theorem 22 and let u tend to the limit $1 - \frac{2i}{n-1}$. This gives: $\widehat{A}_i(n)$ restricted to $SO(n-1, 1)$, contains $\widehat{A}_i(n-1)$ discretely. The rest of the corollary follows by induction. $\square$

Corollary 6. *Suppose that for all n , the **tempered** cohomological representations $A_i(n)$ (i.e. $i = [n/2]$) are not limits of complementary series in the **automorphic dual** of $SO(n, 1)$. Then, for any n , all nontempered cohomological representations of $SO(n, 1)$ are isolated in the automorphic dual.*

Acknowledgments.

T.N.Venkataramana thanks the Department of Mathematics, Cornell University and the Institute for Advanced Study, Princeton for their hospitality in 2007- 2008, where a considerable part of this work was done. He also thanks Peter Sarnak for valuable conversations concerning the material of the paper. He gratefully acknowledges the support of the J.C Bose fellowship for the period 2008-2013.

B. Speh was partially supported by NSF grant DMS 0070561. She would also like to thank the Tata Institute for its hospitality in the winter of 2009 during a crucial stage of this work.

REFERENCES

- [Ba-B] Barbasch and Baldoni Silva, unitary dual for rank one groups, Invent math 1980.
- [Ber] N.Bergeron, Lefschetz Properties for Real and Complex Manifolds, Int. Math. Res. Not., 2003, no.20, 1089-1122.
- [BLS] Burger, M., Li, J-S. and Sarnak, P. Ramanujan Duals and automorphic spectrum, Bull.Amer.Math.Soc. (N.S.)**26** (1992), 253-257.
- [Bu-Sa] Burger, M. and Sarnak, P. Ramanujan Duals II. Invent.Math.**106** (1991) 1-11.
- [Clo] Clozel, L. Démonstration de la conjecture τ . Invent. Math.**151** (2003), 297-328.
- [GW] Goodman and Wallach, Representations and Invariants of the Classical Groups, Encyclopedia of Mathematics and Its Applications, Cambridge University Press, 1998.
- [GV] Graev and Vershik, The Structure of Complementary Series of $O(n, 1)$ and $U(n, 1)$. Russian Math. Surveys, **61**, (2006), **no. 5**, 799-884.
- [Sel] Selberg, A. On the Estimation of Fourier Coefficients of Modular Forms, Proc. Symp. Pure Math. Volume **8**, AMS, Rhode Island, (1965), 1-15.

E-mail address: `venky@math.tifr.res.in`